



**Press release:** Embargoed until **24 July 2019** 18:00 (UK time) / 13:00 (US Eastern time)

## Industrialised fishing overlaps threatened shark hotspots worldwide

- Major high seas fishing activities are centred on ecologically important shark hotspots worldwide
- North Atlantic blue sharks and shortfin mako – the fastest shark in the sea – have on average 76% and 62% of their space use, respectively, overlapped by longlines each month
- Even internationally protected species such as great white and porbeagle sharks at risk
- Danger of accelerating decline in populations and disappearance of hotspots
- Results could provide a ‘blueprint’ for use in deciding where to place large-scale marine protected areas (MPAs)

Even the remotest parts of the ocean appear to offer highly migratory sharks little refuge from industrialised fishing fleets, a ground-breaking study published in the journal *Nature* reports.

Large sharks inhabiting the open ocean account for over half of all identified shark catch globally in target fisheries or as bycatch. Regional declines in abundance of some populations such as shortfin mako shark – the fastest shark in the sea – have led to widespread calls for catch limits in the High Seas (areas beyond national jurisdiction; ABNJ) where there is currently little or no management for sharks.

But precisely where in the vast expanse of the oceans sharks aggregate and how much fishing takes place in those chosen habitats remains poorly known globally, even though it will be crucial to selecting sites to conserve sharks.

Now, an international team of over 150 scientists from 26 countries have tackled this knowledge gap by collating movement data from nearly 2,000 sharks tracked with satellite transmitter tags.

Researchers mapped shark positions and revealed ‘hotspots’ of space use in unprecedented detail. They found multi-species pelagic shark hotspots were mostly located in frontal zones, boundaries in the sea between different water masses that are highly productive and food-rich.

They then calculated how much the hotspots were overlapped by global fleets of large, longline fishing vessels – the type of fishing gear that catches most pelagic sharks – with each vessel capable of deploying 100 km long lines bearing 1200 baited hooks on a daily basis.

Strikingly, they found 24% of the mean monthly space used by sharks globally falls under the footprint of pelagic longline fisheries. For commercially exploited sharks such as North Atlantic blue and shortfin makos sharks overlap was much higher, with on average 76% and 62% of their space use, respectively, overlapping with longlines each month. Even internationally protected species



such as great white and porbeagle sharks had overlap values exceeding 50%.

“Our results show major high seas fishing activities are currently centred on ecologically important shark hotspots worldwide” says Professor David Sims, who led the study as part of the Global Shark Movement Project based at the Marine Biological Association Laboratory in Plymouth, UK.

Equally alarming was that shark hotspots showing high overlap with longline fishing were often also subjected to high fishing effort, a potential ‘double whammy’ for sharks that will result in higher catch rates and potentially accelerate declines in abundance.

“Some shark hotspots were exposed to higher than average fishing effort for as much as half the year” said Dr Nuno Queiroz, a lead researcher in the study from the University of Porto in Portugal.

Given the high fishing effort in hotspots of many species for significant portions of the year, and the very few tracked hotspots free from exploitation, the study reveals exposure risk of sharks to fisheries in the high seas is spatially extensive – stretching across entire ocean-scale population ranges for some species.

The team’s findings indicate large sharks – some of which are already endangered globally – face a future with limited spatial refuge from industrial longline fishing effort.

The researchers propose that the detailed maps of shark hotspots and exposure to longline fishing effort they have produced can provide a ‘blueprint’ for use in deciding where to place large-scale marine protected areas (MPAs) aimed at conserving sharks, in addition to the need for strict quotas to reduce catches elsewhere.

“Some of the shark hotspots we studied may not be there in as little as a few years’ time if management measures are not put in place now to conserve the sharks and the habitats on which they depend” Sims says.

The research is published online in *Nature* on 24<sup>th</sup> July 2019.

## Ends

### **Notes for editors:**

1. Contacts

Corresponding author of the research: Prof David Sims, MBA Laboratory, Plymouth, UK.

Landline: +44 (0)1752 426487. Mobile: +44 (0)7848 028456. Email: [dws@mba.ac.uk](mailto:dws@mba.ac.uk) Twitter: @TheSimsLab

Guy Baker, Communications Co-ordinator

Marine Biological Association (MBA), The Laboratory, Citadel Hill, Plymouth, Devon, PL1 2PB, UK [www.mba.ac.uk](http://www.mba.ac.uk)

Landline: +44 (0)1752 426239 Mobile: +44 (0)7876 831267 Email: [guba@mba.ac.uk](mailto:guba@mba.ac.uk)



3. The full citation to the paper is: Queiroz, N., Humphries, N.E., Couto, A., Vedor, M., da Costa, I., Sequeira, A.M.M., Mucientes, G., Santos, A.M., Abascal, F.J., Abercrombie, D.L., Abrantes, K., Acuña-Marrero, D., Afonso, A.S., Afonso, P., Anders, D., Araujo, G., Arauz, R., Bach, P., Barnett, A., Bernal, D., Berumen, M.L., Bessudo Lion, S., Bezerra, N.P.A., Blaison, A.V., Block, B.A., Bond, M.E., Bradford, R.W., Braun, C.D., Brooks, E.J., Brooks, A., Brown, J., Bruce, B.D., Byrne, M.E., Campana, S.E., Carlisle, A.B., Chapman, D.D., Chapple, T.K., Chisholm, J., Clarke, C.R., Clua, E.G., Cochran, J.E.M., Crochelet, E.C., Dagorn, L., Daly, R., Devia Cortés, D., Doyle, T.K., Drew, M., Duffy, C.A.J., Erikson, T., Espinoza, E., Ferreira, L.C., Ferretti, F., Filmlalter, J.D., Fischer, C.G., Fitzpatrick, R., Fontes, J., Forget, F., Fowler, M., Francis, M.P., Gallagher, A.J., Gennari, E., Goldsworthy, S.D., Gollock, M.J., Green, J.R., Gustafson, J.A., Guttridge, T.L., Guzman, H.M., Hammerschlag, N., Harman, L., Hazin, F.H.V., Heard, M., Hearn, A.R., Holdsworth, J.C., Holmes, B.J., Howey, L.A., Hoyos, M., Hueter, R.E., Hussey, N.E., Huveneers, C., Irion, D.T., Jacoby, D.M.P., Jewell, O.J.D., Johnson, R., Jordan, L.K.B., Jorgensen, S.J., Joyce, W., Keating Daly, C.A., Ketchum, J.T., Klimley, A.P., Kock, A.A., Koen, P., Ladino, F., Lana, F.O., Lea, J.S.E., Llewellyn, F., Lyon, W.S., MacDonnell, A., Macena, B.C.L., Marshall, H., McAllister, J.D., McAuley, R., Meÿer, M.A., Morris, J.J., Nelson, E.R., Papastamatiou, Y.P., Patterson, T.A., Peñaherrera-Palma, C., Pepperell, J.G., Pierce, S.J., Poisson, F., Quintero, L.M., Richardson, A., Rogers, P.J., Rohner, C.A., Rowat, D.R.L., Samoily, M., Semmens, J.M., Sheaves, M., Shillinger, G., Shivji, M., Singh, S., Skomal, G.B., Smale, M.J., Snyders, L.B., Soler, G., Soria, M., Stehfest, K.M., Stevens, J.D., Thorrold, S.R., Tolotti, M.T., Towner, A., Travassos, P., Tyminski, J.P., Vandeperre, F., Vaudo, J.J., Watanabe, Y.Y., Weber, S.B., Wetherbee, B.M., White, T.D., Williams, S., Zárata, P.M., Harcourt, R., Hays, G.C., Meekan, M.G., Thums, M., Irigoien, X., Eguiluz, V.M., Duarte, C.M., Sousa, L.L., Simpson, S.J., Southall, E.J. & Sims, D.W. (2019) Global spatial risk assessment of sharks under the footprint of fisheries. *Nature*, doi: 10.1038/s41586-019-1444-4
4. The analysis was funded in part by the [UK Natural Environment Research Council](#) (NERC) and Fundação para a Ciência e a Tecnologia (FCT). Field research was supported by many funders listed at the end of the research paper's Supplementary Information.
5. A full list of participating institutions, organisations and research group leaders is given here: [www.globalsharkmovement.org/people](http://www.globalsharkmovement.org/people)

6. The Global Shark Movement Project ([www.globalsharkmovement.org](http://www.globalsharkmovement.org)) is a collaborative scientific research project based at the Laboratory of the Marine Biological Association in Plymouth, UK ([www.mba.ac.uk](http://www.mba.ac.uk)), which aims to advance scientific knowledge of shark behaviour, ecology, conservation and fisheries science that can be used to inform improved management of threatened sharks and ocean biodiversity.



Tens of millions of pelagic sharks are harvested each year by high seas fisheries but with little or no management for the majority of species. Sharks are particularly susceptible to the effects of human exploitation due to slow growth rates, late age at maturity and low fecundity, traits making them comparable to marine mammals in terms of vulnerability. Global data on pelagic shark habitat hotspots, spatial patterns of vulnerability in relation to fisheries, and how sharks respond to changing environment are lacking, precluding understanding where in the global ocean conservation needs to be focused. GSMP aims to close this knowledge gap.

7. The Global Fishing Watch ([www.globalfishingwatch.org](http://www.globalfishingwatch.org)) and Tagging of Pacific Predators ([www.gtopp.org](http://www.gtopp.org)) programs kindly made data freely available in support of this research.



**The Marine Biological Association (MBA)** is a professional body for marine scientists with some 1,700 members world-wide. Since 1884 the MBA has established itself as a leading marine biological research organization contributing to the work of several Nobel Laureates and over 170 Fellows of the Royal Society. In 2013, the MBA was awarded a Royal Charter in recognition of its long and eminent history and its status within the field of marine biology. The award strengthens the Association's role in promoting marine biology as a discipline and in representing the interests of the marine biological community. The Association is based at its world-class marine research laboratory in Plymouth.

[www.mba.ac.uk](http://www.mba.ac.uk) Twitter: @thembauk

